

OR-1. 2-ARYL AND 2-THIENYLBENZAZINES WITH LUMINESCENT PROPERTIES

T. N. Moshkina¹, E. V. Nosova^{1,2}, G. N. Lipunova², V. N. Charushin^{1,2}


¹Ural Federal University of the first President of Russia B. N. Yeltsin,
Mira St., 19, Yekaterinburg, 620002, Russia

²I. Ya. Postovsky Institute of Organic Synthesis UB RAS,
S. Kovalevskoy/Akademicheskaya St., 20/22, Yekaterinburg, 620990, Russia

E-mail: tan.moshckina@yandex.ru

The benzazine core due to electron withdrawing properties is considered as perspective fragment for construction of π -conjugated systems with push–pull architecture [1, 2]. 2-Hydroxybenzazines can be used as N, O-bidentate ligands for creation of boron chelate complexes possessing promising luminescence properties [3, 4].

The novel push-pull systems (**1–3**) were obtained via Suzuki and Sonogashira Pd-catalyzed cross-coupling reactions. The photophysical properties of compounds **1–3** have been investigated in two solvents, and effect of protonation has been studied [5–8]. Some quinazoline derivatives possess intensive luminescence in toluene (up to 82 %).



R = morpholin-1-yl, CN, OH; **R**¹ = Ar, 2-thienyl, Ph–Ca≡C–, 4-MeOC₆H₄–Ca≡C–; **Ar** = Ph, 4-methoxythienyl, 3,4,5-trimethoxyphenyl, 4-(diethylamino)phenyl, 4-(diphenylamino)phenyl, 4-(9H-carbazol-9-yl)phenyl, **Ar**¹ = Ph, 2-fluorenyl, cymantrenyl; **R**² = F, Ph; **R**³ = H, t-Bu; **X** = H, Cl

Coordination of N, O-ligands with boron trifluoride etherate led to BF₂ complexes (**4–6**) under rather mild conditions. The correlation between compounds structure and the fluorescence intensity, absorption and emission maximum in solution has been shown [9]. Some of samples demonstrated strong blue-green luminescence in the solid state.

References

1. Functionalized Quinazolines and Pyrimidines for Optoelectronic Materials / G. N. Lipunova [et al.] // Curr. Org. Synth. 2018. Vol. 15, № 6. P. 793–814.
2. Protonable pyrimidine derivative for white light emission / S. Achelle [et al.] // RSC Adv. 2015. Vol. 5, № 130. P. 107396–107399.
3. Photophysical Properties of Phenacylphenanthridine Difluoroboranyls: Effect of Substituent and Double Benzannulation / A. M. Grabarz [et al.] // J. Org. Chem. 2017. Vol. 82, № 3. P. 1529–1537.

4. Synthesis and antitumour activity of 4-aminoquinazoline derivatives / G. N. Lipunova [et al.] // Russ. Chem. Rev. 2016. Vol. 85, № 7. P. 759–793.
5. Synthesis and Photophysical Studies of 2-(Thiophen-2-yl)-4-(morpholin-4-yl)quinazoline Derivatives / E. V. Nosova [et al.] // European J. Org. Chem. 2016. Vol. 2016, № 16. P. 2876–2881.
6. Synthesis and photophysical studies of novel 2-[5-(4-diethylaminophenyl)thiophen-2-yl]quinazoline derivatives / E. V. Nosova [et al.] // Mendeleev Commun. 2018. Vol. 28, № 1. P. 14–16.
7. Synthesis and photophysical studies of novel 4-aryl substituted 2-phenyl-, 2-(fluoren-2-yl)- and 2-cyanotrenylquinazolines / E. V. Nosova [et al.] // Mendeleev Commun. 2018. Vol. 28, № 1. P. 33–35.
8. New 2,3-Bis(5-arylthiophen-2-yl)quinoxaline Derivatives: Synthesis and Photophysical Properties / T. N. Moshkina [et al.] // Asian J. Org. Chem. 2018. Vol. 7, № 6. P. 1080–1084.
9. Synthesis, structure and photoluminescent properties of BF₂ and BPh₂ complexes with N,O-benzazine ligands / E. V. Nosova [et al.] // J. Fluor. Chem. 2015. Vol. 175. P. 145–151.

This work was supported by the Russian Foundation for Basic Research (grant № 18-03-00112).